

## **Considerations in performing endoscopy during the COVID-19 pandemic**

Roy Soetikno, MD, MS<sup>1</sup>, Anthony YB Teoh, MBChB<sup>2</sup>, Tonya Kaltenbach, MD, MS<sup>1,3</sup>,  
James Y.W. Lau, MD<sup>2</sup>, Ravishankar Asokkumar, MBBS<sup>4,5</sup>, Patricia Cabral-Prodigalidad,  
MD<sup>6</sup>, Amandeep Shergill, MD, MS<sup>1,3</sup>

1. Division of Gastroenterology and Hepatology, San Francisco Veterans Affairs Medical Center, San Francisco, CA, USA
2. Department of Surgery, Chinese University of Hong Kong, Shatin, Hong Kong
3. Department of Medicine, University of California San Francisco, CA, USA
4. Department of Gastroenterology and Hepatology, Singapore General Hospital, Singapore
5. HM Sanchinarro University Hospital, Madrid, Spain.
6. Institute of Digestive and Liver Diseases, St. Luke's Medical Center-Global City, Taguig, Philippines

### **Send correspondence to:**

Roy Soetikno, MD, MS

San Francisco VA, San Francisco, CA

[soetikno@earthlink.net](mailto:soetikno@earthlink.net)

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

**Considerations in performing endoscopy during the COVID-19 pandemic**

Roy Soetikno, MD, MS<sup>1</sup>, Anthony YB Teoh, MBChB<sup>2</sup>, Tonya Kaltenbach, MD, MS<sup>1,3</sup>,  
James YW Lau, MD<sup>2</sup>, Ravishankar Asokkumar, MBBS<sup>4,5</sup>, Patricia Cabral-Prodigalidad,  
MD<sup>6</sup>, Amandeep Shergill, MD, MS<sup>1,3</sup>

1. Division of Gastroenterology and Hepatology, San Francisco Veterans Affairs Medical Center, San Francisco, CA, USA
2. Department of Surgery, Chinese University of Hong Kong, Shatin, Hong Kong
3. Department of Medicine, University of California San Francisco, CA, USA
4. Department of Gastroenterology and Hepatology, Singapore General Hospital, Singapore
5. HM Sanchinarro University Hospital, Madrid, Spain.
6. Institute of Digestive and Liver Diseases, St. Luke's Medical Center-Global City, Taguig, Philippines

**Send correspondence to:**

Roy Soetikno, MD, MS  
San Francisco VA, San Francisco, CA  
[soetikno@earthlink.net](mailto:soetikno@earthlink.net)

**Acronyms:**

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

**SARS-CoV-2:** Severe acute respiratory syndrome coronavirus 2

**COVID-19:** Coronavirus Disease 2019

**HCP:** Health Care Providers

**PPE:** Personal Protective Equipment

**PAPR:** Powered Air Purifying Respirator

**AGP:** Aerosol Generating Procedure

**WHO:** World Health Organization

**CDC:** Centers for Disease Control and Prevention

**PUI:** Person under investigation

**BSL:** Biosafety Level

1  
2  
3  
4 **Introduction**  
5

6  
7 Based on experiences and the literature, our objective is to provide practical suggestions  
8  
9 for performing endoscopy in the setting of COVID-19 pandemic.  
10

11  
12  
13  
14 SARS-CoV-2/Novel Coronavirus-19 (COVID-19) has become a global pandemic.  
15  
16 Human-to-human transmission occurs through respiratory secretions, aerosols, feces,  
17  
18 and contaminated environmental surfaces.<sup>1,2</sup> Transmission can occur in both  
19  
20 symptomatic and asymptomatic individuals.<sup>3</sup> Viable virus particles can be detected in  
21  
22 aerosols up to 3 hours after aerosolization and up to 3 days on surfaces.<sup>4</sup> A recent  
23  
24 publication suggests that undocumented infections were the source of a substantial  
25  
26 majority of documented cases.<sup>5</sup> The risk of infection to healthcare workers is significant:  
27  
28 in one of the earliest documentations of infection in Wuhan, 29% of patients (40 out of  
29  
30 138) were healthcare workers.<sup>6</sup> It is unknown how much of the risk was related to the  
31  
32 direct care of infected patients or to the inadequate use of personal protective equipment  
33  
34 (PPE).  
35  
36  
37  
38  
39  
40  
41  
42

43 When performing endoscopy, it seems inevitable that healthcare providers (HCP) will be  
44  
45 exposed to either respiratory or gastrointestinal fluids from patients. Thus, adequate  
46  
47 protection of HCP is now critical. The World Endoscopy Organization has recently  
48  
49 released a recommendation on infection prevention and control in digestive endoscopy  
50  
51 based on experiences from China.<sup>7</sup> Similarly, an Italian group has provided  
52  
53 recommendations regarding the performance of endoscopy during the COVID-19  
54  
55 outbreak.<sup>8</sup>  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4  
5  
6  
7 Based on their experiences with a similar coronavirus, which caused SARS 17 years ago,  
8  
9 Hong Kong adopted measures similar to those aforementioned immediately upon the first  
10  
11 news of COVID-19 outbreak in late January.<sup>9</sup> With numbers of COVID-19 cases  
12  
13 continuing to rise in North America and Europe, we aim to provide practical suggestions  
14  
15 to potentially avoid the transmissions of COVID-19 in the endoscopy unit.  
16  
17  
18  
19  
20

### 21 **Potential mode of transmissions of SARS-CoV-2 during endoscopy**

22

23 The virus characteristics and its transmission make endoscopy a potential route for  
24  
25 infection. Possible routes of SARS-CoV-2 transmission include (1) person-to-person, (2)  
26  
27 respiratory droplets, (3) aerosols generated during endoscopy, and (4) contact with  
28  
29 contaminated surroundings and body fluids.<sup>1,10</sup> Additional care must be instituted when  
30  
31 handling blood samples or specimens because the virus has been detected in the blood  
32  
33 of COVID-19 patients. Lei Pan et al<sup>10</sup> demonstrated that 48.5% of the patients presented  
34  
35 with GI symptoms, including anorexia (83.8%), diarrhea (29.3%), and vomiting (0.8%),  
36  
37 with the severity increasing as the disease progressed. With the detection of the virus in  
38  
39 the feces, the Centers for Disease Control and Prevention (CDC) has suggested the use  
40  
41 of separate bathrooms in cases of suspected COVID-19.<sup>11</sup> In line with these  
42  
43 recommendations, extensive precautions need to be adopted to avoid potential oral-fecal  
44  
45 transmission.  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4 Importantly, staff with a travel history to COVID-19–affected areas or a history of exposure  
5  
6 to COVID-19–affected individuals should first self-quarantine for 14 days, to eliminate risk  
7  
8 of transmission.  
9

10  
11  
12  
13  
14 *Is endoscopy an aerosol-generating procedure?*

15 All endoscopic procedures should be considered aerosol-generating procedures (AGP).  
16  
17 Coughing and retching can occur during upper endoscopy, generating aerosols. Likewise,  
18  
19 patients undergoing colonoscopy may pass flatus, which is also known to disseminate  
20  
21 bacteria to nearby surroundings.<sup>12</sup> A prospective study has demonstrated unrecognized  
22  
23 endoscopist exposure to infectious particles during GI procedures.<sup>13</sup> Recently, the World  
24  
25 Health Organization (WHO) has published an extensive guideline on the rational use of  
26  
27 personal protective equipment (PPE) for COVID-19 and provided specific instructions for  
28  
29 healthcare workers performing AGP on patients with COVID-19.<sup>14</sup> These include the use  
30  
31 of a respirator (N95, FFP2 standard, or equivalent), gown, gloves, eye protection, and  
32  
33 apron although aprons are not usually not used in the United States. Their use should be  
34  
35 immediately and strictly adopted in practice, if at all possible.<sup>13</sup>  
36  
37  
38  
39  
40  
41  
42  
43  
44

45 *Surface contamination during endoscopy*

46  
47 Patient-contaminated fluids often splatter when inserting or removing an accessory from  
48  
49 the endoscope’s working channel, adjusting the air/water button, retrieving tissue from a  
50  
51 biopsy bottle, and while performing precleaning. Patients’ saliva can contaminate the  
52  
53 pillow or the bed, and stool mixed with water often drips to the bed during colonoscopy.  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4 *Contamination of the room used by patients with COVID-19*

5  
6  
7 Extensive environmental contamination can occur even from patients with mild COVID-  
8  
9 19 upper respiratory symptoms. Ong and colleagues<sup>15</sup> detected positive SARS-CoV-2  
10  
11 samples in various locations around the patient's room, including the patient's bed, sink,  
12  
13 bathroom, light switches, and doors. In addition, positive samples were found on the  
14  
15 shoes and stethoscope of staff exiting the patient's room. However, there was no  
16  
17 contamination in the anteroom or corridor outside the room. The study illustrates the  
18  
19 significant extent of contamination by patients with SARS-CoV-2 through respiratory  
20  
21 droplets and fecal shedding.<sup>16</sup>  
22  
23  
24  
25  
26  
27

28 **Our Challenge**

29  
30  
31 The goal is to attain zero percent infection rates among HCP while providing essential  
32  
33 services to patients. For the GI community, the key element will be to prevent exposure  
34  
35 during any endoscopic procedure. As the outbreaks continue to occur, masks and  
36  
37 personal protective equipment (PPE) may become scarce in quantity. An early inventory  
38  
39 of what is available to the institute is essential to formulate a plan for PPE usage.  
40  
41 Conservation of PPE is important and should be planned.  
42  
43  
44  
45  
46  
47

48 **Suggested general measures**

49  
50 ***Management***

- 51  
52  
53 1. Prepare. Plan. Test. Practice. Repeat. Ready the team. Being well prepared is the  
54  
55 best we can do to reach our zero-contamination goal.  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

- 1
- 2
- 3
- 4 2. Staff management is an integral part of performing endoscopy during the COVID
- 5 pandemic.
- 6
- 7
- 8
- 9 3. Establishment of a rapid response communication channel using smart phone
- 10 apps, email, and video conferences to distribute information across the entire unit.
- 11
- 12
- 13 4. Regular updates with administrators, infectious control team, doctors, and nurses
- 14 to stay updated on the development of the infection and discuss a unified plan.
- 15
- 16
- 17
- 18
- 19
- 20

### 21 ***Preparation***

- 22
- 23 1. Ensure performance of fit testing for N95 respiratory masks for all HCP. During the
- 24 course of the outbreak, some masks may run out of stock and HCP will need to
- 25 plan for alternatives. Protection, however, may be achievable even without N95
- 26 through the use of medical masks.<sup>17</sup> Note that as an AGP, endoscopy of
- 27 PUI/COVID patients requires the use of respiratory protection. The powered air
- 28 purifying respirator (PAPR) is a desirable alternative that does not require fit testing
- 29 and can be used by employees with facial hair who would otherwise not achieve a
- 30 good seal with the N95. Most units, however, are not stocked to have an adequate
- 31 supply of PAPR.
- 32
- 33
- 34
- 35
- 36
- 37
- 38
- 39
- 40
- 41
- 42
- 43
- 44
- 45 5. Familiarize staff with the correct method of hand hygiene. An excellent review has
- 46 been published.<sup>18</sup> Compliance with correct hand hygiene practices is low; thus, a
- 47 practice, review, and compliance check is necessary.
- 48
- 49
- 50
- 51
- 52
- 53 6. Follow the WHO recommendations for PPE (Table 1).<sup>14</sup> Familiarize staff with the
- 54 correct sequence of gowning up (donning) and down (doffing) through teaching
- 55 videos and diagrams (Table 1)<sup>19</sup>. Note that there is poor correlation between self-
- 56
- 57
- 58
- 59
- 60
- 61
- 62
- 63
- 64
- 65



1  
2  
3  
4 perceived proficiency in PPE use and its appropriate use.<sup>17</sup> Repetitive training and  
5 demonstrated competency are necessary. Use a buddy system, where another  
6 colleague observes the gown up and down procedures to advise on any breach of  
7 protocol (Table 2). Inform to conserve the use of masks and PPE.

- 8  
9  
10  
11  
12  
13  
14 7. Setting up a reception bay to screen and stratify risk of COVID-19 infection of the  
15 patient before allowing them to enter the waiting area alongside other patients. In  
16 the setting of substantial community spread, make efforts to separate all patients  
17 by approximately 6 feet.  
18  
19  
20  
21  
22  
23 8. Set up a designated procedure and recovery room for suspected (PUI) and  
24 COVID-19 positive patients. The endoscopic procedures should be performed in  
25 an Airborne Infection Isolation Room that adheres to Level 3 biosafety  
26 requirements. Consider alternative sites with enhanced prevention capabilities for  
27 performing procedures if an isolation room is not available.  
28  
29  
30  
31  
32  
33 9. Set up a designated area for donning PPE that is easily accessible and near the  
34 room. Doffing of PPE ideally occurs in an anteroom or a doffing area that is  
35 separate from the procedure room.<sup>20</sup>  
36  
37  
38  
39  
40  
41  
42  
43 10 Equally important, staff should take additional precautions to prevent  
44 contamination among providers. Work at individual working stations using a  
45 designated phone, computer, and chair, and stay at least 6 feet from any other  
46 coworkers while at work to the extent possible, recognizing that this will be difficult  
47 in some situations. Avoid sharing workstation items and equipment. Wipe  
48 workstations before and after use with virucide, following instructions on the  
49 virucide exactly as recommended.  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

- 1  
2  
3  
4  
5  
6  
7 11 Create a workflow to provide a clear job description and designation of authority  
8  
9 with backup plans. Separate the workflow to minimize cross-contamination. For  
10  
11 example, consider dividing the clinical workforce into 2 teams, alternating roles at  
12  
13 predefined intervals (such as weekly). One team is on-site and providing direct  
14  
15 clinical care. The second team is coordinating clinical care off-site, minimizing risk  
16  
17 of exposure and providing back-up coverage if an on-site provider were to become  
18  
19 ill or require quarantine.  
20  
21  
22  
23 12 In the staff lounge/eating area, we allow sitting in one direction, thus preventing  
24  
25 infection from face-to-face transmission.  
26  
27  
28 13 The bathroom is a potential site of transmission. Ideally patient and staff bathrooms  
29  
30 are separated and disinfected frequently.  
31  
32  
33  
34  
35

### ***Indications for procedures***

36  
37  
38 In the epidemic area, the indications include management of upper gastrointestinal  
39  
40 bleeding, acute cholangitis, foreign body, and obstructions.<sup>7</sup> Care (initial diagnosis,  
41  
42 biopsy, staging, palliation of biliary and luminal obstruction) of cancer patients may also  
43  
44 be considered urgent. Reschedule nonurgent endoscopy services. This measure is aimed  
45  
46 at reducing the risk of spreading infection from asymptomatic patients, reducing the risk  
47  
48 of cross-infection among patients, reducing use of PPE, and reducing unnecessary  
49  
50 admissions to free up hospital resources.  
51  
52  
53  
54  
55  
56

### ***In Practice***

1  
2  
3  
4 *In the setting of substantial community spread of COVID-19:*  
5

- 6  
7 1. Require all staff to have daily measurements of temperature before starting work.  
8 All febrile staff should not be allowed to work, and they should be evaluated  
9 according to local protocols to screen for potential COVID-19 infections.  
10
- 11 2. CDC mitigation strategies in the setting of substantial community spread include  
12 requiring all HCP to wear a face mask when in the facility, depending on supply.<sup>21</sup>  
13  
14 There is a high viral load in the upper respiratory tract, and there is a significant  
15 potential for asymptomatic persons to shed and transmit virus.<sup>22</sup> Data showing the  
16 prolonged stability of the virus on surfaces may have significant potential  
17 implications for use of staff' PPE in the general area.<sup>23</sup>  
18
- 19 3. Require staff to perform work using individual stations: use the same phone,  
20 computer, and chair. Do not share. Do not answer phones elsewhere other than in  
21 your own station and disinfect your working space regularly.  
22
- 23 4. Limit the number of HCP in the endoscopy suite to those essential for performance  
24 of the procedures (see below regarding trainees). Off-duty workers should stay at  
25 home as much as possible.  
26
- 27 5. For HCP directly involved in the procedures, use the hospital-issued scrubs and  
28 dedicated endoscopy shoes. Leave these at work.  
29
- 30 6. Although these continue to evolve, current COVID-19 screening guidelines include  
31 assessing patient's symptoms (such as fever and/or symptoms of acute respiratory  
32 illness) and potential contact with a suspected or laboratory-confirmed COVID-19  
33 patient. The decision to quarantine should be made at that time (Fig. 1).  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

- 1  
2  
3  
4 7. With the availability of RNA testing against COVID, point-of-care testing in patients  
5 presenting for endoscopy may facilitate more accurate risk stratification.  
6  
7  
8  
9

10  
11 *Before the procedure:*  
12

13  
14 *Outpatients*  
15

- 16 1. Screen for symptoms, signs, and exposure to SARS-CoV-2 (contact and travel  
17 history). Measure their temperature to risk stratify (Fig. 1).  
18  
19 2. Test all suspected patients for COVID-19 whenever possible using RT-PCR.<sup>24</sup> If  
20 possible, wait until the test results have been received before proceeding.  
21  
22 3. Suspected or confirmed patients should be provided a mask while being triaged,  
23 and should be isolated or separated from other patients by at least 6 feet.  
24 Alternately, they should be placed in a negative pressure room.  
25  
26 4. Patients should be advised to minimize movements while waiting for the procedure  
27 to minimize facility contamination.  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37

38 *In-patients*  
39

- 40 1. Evaluate for COVID-19 status and reassess for symptoms suspicious for COVID-  
41 19 in all patients referred for endoscopy and triage accordingly.  
42  
43 2. PUI and COVID patients should be provided a mask while awaiting the procedure  
44 and should be stationed away from other patients as detailed above. Designated  
45 transportation corridors or lift/elevator should be used to transfer patients to the  
46 endoscopy unit.  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57

58 *During the procedure for PUI/ COVID-19 positive patients:*  
59  
60  
61  
62  
63  
64  
65

- 1
- 2
- 3
- 4 1. Include verification of the patient's status for COVID-19 in the Time Out protocol.
- 5
- 6 2. Ensure that a PPE supply is available before entering the procedure room.
- 7
- 8
- 9 3. Wash your hands according to the recommended hand-washing method.
- 10
- 11 4. Follow the CDC's recommendations for performing AGP: wear a respirator (N95,
- 12 FFP2 standard, or equivalent); impermeable gown; gloves; apron; and eye
- 13 protection (Fig. 2).
- 14
- 15 5. Follow the CDC sequence for putting on the PPE. Remove all personal items, such
- 16 as jewelry, pagers, and badges. Use the buddy system to confirm that the PPE is
- 17 correctly in place for those who are not familiar with gowning up and down.
- 18
- 19 6. Consider boot covers during ERCP.
- 20
- 21 7. If the patient requires tracheal intubation for the procedure, only the
- 22 anesthesiologist and his/her assistant stay in the room during intubation. The
- 23 remaining team stays outside.
- 24
- 25 8. After the procedure is completed, follow the CDC's recommendations for taking off
- 26 PPE. Use a buddy system to observe for any breach. If there is, use an alcohol
- 27 spray to decontaminate the potentially touched area(s).
- 28
- 29 9. Wash your hands according to the recommended hand-washing method. Confirm
- 30 proficiency.<sup>18</sup>
- 31
- 32 10. Other HCP in the clean area can complete the procedure report, thus potentially
- 33 avoiding contaminations.<sup>7</sup>
- 34
- 35
- 36
- 37
- 38
- 39
- 40
- 41
- 42
- 43
- 44
- 45
- 46
- 47
- 48
- 49
- 50
- 51
- 52
- 53
- 54
- 55
- 56
- 57
- 58
- 59
- 60
- 61
- 62
- 63
- 64
- 65

*After the procedure, all patients:*

- 1  
2  
3  
4 1. Staff: After performing endoscopy in a person under investigation (PUI)/COVID-19  
5  
6 positive patient, shower before leaving the hospital.  
7  
8  
9 2. Patients:  
10  
11 a. Provide patients with a suitable PPE, depending on their risk status, while  
12 waiting in the recovery area.  
13  
14 b. Toilet flush is known to generate bioaerosols (“Toilet plume”) and may  
15 transmit infection. Advise patients to flush the toilets after use with the lids  
16 closed.  
17  
18 c. Contact asymptomatic patients within 14 days to assess their progress after  
19 procedure.  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30

31 *Disinfection management:*

32  
33 Reprocessing reusable medical equipment. We are not aware of a change in the RME  
34 protocol. Note that the most significant HCP contamination occurs during precleaning of  
35 the endoscope in the procedure room due to splashing from the air/water button. Follow  
36 the protocol to turn off the processor when replacing the air/water button with the credit  
37 card button.  
38  
39  
40  
41  
42  
43  
44  
45

46  
47 SARS-CoV-2 is deactivated by commonly used disinfectants such as alcohol or chlorine-  
48 based solutions. The CDC cleaning and disinfection recommendation can be adopted.  
49  
50 Please see Table 2 for the recommended attire for the personnel cleaning the unit.  
51  
52 Personnel cleaning the endoscopy unit must also undergo repeated practice and have  
53 their proficiency documented.  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4  
5  
6  
7 Special considerations: trainee involvement  
8

9 Trainees are an integral part of most academic endoscopic units. With the potential surge  
10 in COVID-19 infection, the role of a trainee in endoscopy procedures requires re-  
11 evaluation. Because there is too much uncertainty with regard to its transmissible  
12 potential and associated morbidity and mortality, we recommend the following plan of  
13 actions in managing trainees during endoscopy:  
14  
15  
16  
17  
18  
19  
20

- 21 1. They master the prevention of transmission described previously through repeated  
22 practice and documented proficiency.  
23
- 24 2. Fellows' involvement increases procedure time, and thus increases the potential  
25 for exposure. Our practice is to preserve critical resources and minimize the risk  
26 of exposure; thus, we limit trainees' involvement during endoscopic procedures.<sup>25</sup>  
27  
28 As board-certified internists, however, fellows may provide essential physician  
29 support in a time of crisis, such as during a surge. They may contribute to the  
30 COVID-19 management workforce.  
31  
32
- 33 3. At many institutions, fellows cover multiple clinical sites as part of their on-call  
34 duties or for ACGME required continuity clinics. In the absence of point-of-care  
35 testing, we suggest that fellows be stationed at one hospital to avoid inadvertent  
36 spread of infection across multiple sites.  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52

53 Our guidance is based on our practical experience, observations, and published literature.  
54

55 Note that present understanding of SARS-CoV-2, however, is still rapidly evolving.  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

The success of preventing endoscopy unit transmission of SARS-CoV-2 is contingent upon the compliance of every member of the team. We must cooperate and collaborate in order to adhere to the prevention steps the best we can and prevent transmissions.

**Figure legends**









**Figure 1.** A schematic representation of the workflow for managing suspected (PUI) or COVID-19 patients in an endoscopy unit. Note that this workflow is still evolving. The availability of testing kits would likely change it.

**Figure 2.** Typical PPE used in our unit in Hong Kong (Image courtesy of Dr Anthony Teoh, Hong Kong, China). Designated shoes to be worn in the endoscopy unit only.



**Tables**

Table 1. Recommended protocol for putting on and removing personal protective equipment (PPE). Adapted from CDC guidelines

<p><b>How to put on PPE</b></p>	<p><b>1. Gown</b> Fully cover torso from neck to knees, arms, to end of wrists and wrap around the back</p> 	<p><b>2. Mask</b> Secure ties or elastic bands at the middle of head and neck. Fit flexible band to nose bridge. Fit snug to face and below chin. Fit- check respirator</p> 	<p><b>3. Goggles</b> Place over face and eyes to adjust fit.</p> 	<p><b>4. Gloves</b> Extend to cover the wrist of isolation gown</p> 
<p><b>How to remove PPE (Example 1)</b></p>	<p><b>1. Gloves</b> Grasp palm area of the other gloved hand and peel off first glove. Hold removed glove in gloved hand. Slide fingers under the glove at the wrist and peel off the second glove over the first.</p> 	<p><b>2. Goggles</b> Lift headband or earpiece from the back to remove goggles or face shield.</p> 	<p><b>3. Gown</b> Unfasten gown ties while ensuring the sleeves do not contact your body. Pull the gown away from the neck by touching the inside of the gown only. Turn inside out and roll into a bundle to discard.</p> 	<p><b>4. Mask</b> Grasp bottom and top ties of the mask. Remove ties without contacting the front of the mask.</p> 

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65




<p><b>How to remove PPE (Example 2)</b></p>	<p><b>1. Gown and gloves</b> Grasp gown in the front and pull away from your body so the ties break. Touch outside of the gown only with gloved hands. While removing the gown, roll it inside-out into a bundle and peel your gloves off at the same time.</p> 	<p><b>2. Goggles</b> Lift headband or earpiece from the back to remove goggles or face shield.</p> 	<p><b>3. Mask</b> Grasp bottom and top ties of the mask. Remove ties without contacting the front of the mask.</p> 
---	---	---	--

Table 2. Recommended personal protective equipment (PPE) to be used in the context of COVID-19 disease, according to the setting, personnel and type of activity.

(\*) Use the buddy system to confirm that PPE is correctly in place. Adapted from WHO and CDC guidelines.

Setting	Target personnel or patients	Activity	Type of PPE or procedure
<b>Healthcare facilities, Inpatient facilities</b>			
Patient room	Healthcare workers	Providing direct care to COVID-19 patients.	*Medical mask, Gown, Gloves Eye protection (goggles or face shield).
	*Use buddy system to prevent protocol breach	Aerosol-generating procedures performed on COVID-19 patients.	*Respirator N95 or FFP2 standard, or equivalent. Gown, Gloves, Eye protection, Apron
	Cleaners	Entering the room of COVID-19 patients.	Medical mask, Gown, Heavy duty gloves Eye protection (if risk of splash from organic material or chemicals). Boots or closed work shoes
	Visitors	Entering the room of a COVID-19 patient	Medical mask, Gown, Gloves
Other areas of patient transit (eg, wards, corridors).	All staff, including healthcare workers.	Any activity that does not involve contact with COVID-19 patients.	Medical mask

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

Triage	Healthcare workers	Preliminary screening not involving direct contact.	Maintain spatial distance of at least 1 meter Medical mask required <sup>21</sup>
	Patients with respiratory symptoms.	Any	Maintain spatial distance of at least 1 meter Provide a medical mask if tolerated by patients
	Patients without respiratory symptoms.	Any	No PPE required
<b>Outpatient facilities</b>			
Consultation room	Healthcare workers	Physical examination of patients with respiratory symptoms.	*Medical mask, Gown, Gloves, Eye protection
	Healthcare workers	Physical examination of patients without respiratory symptoms.	PPE according to standard precautions and risk assessment.
	Patients with respiratory symptoms.	Any	Provide a medical mask if tolerated.
	Patients without respiratory symptoms.	Any	No PPE required
	Cleaners	After and between consultations with patients with respiratory symptoms.	Medical mask, Gown, Heavy duty gloves Eye protection (if risk of splash from organic material or chemicals), Boots or closed work shoes
Waiting room	Patients with respiratory symptoms.	Any	Provide a medical mask if tolerated. Immediately move the patient to an isolation room or separate area away from others; if this is not

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

			feasible, ensure spatial distance of at least 1 meter from other patients.
	Patients without respiratory symptoms.	Any	No PPE required
Triage	Healthcare workers	Preliminary screening not involving direct contact.	Maintain spatial distance of at least 1 meter Medical mask required. <sup>21</sup>
	Patients with respiratory symptoms.	Any	Maintain spatial distance of at least 1 meter Provide medical mask if tolerated.
	Patients without respiratory symptoms.	Any	No PPE required

## References

1. Rio C del, Malani PN. COVID-19—New Insights on a Rapidly Changing Epidemic. *JAMA*. February 2020. doi:10.1001/jama.2020.3072
2. Xiao F, Tang M, Zheng X, Liu Y, Li X, Shan H. Evidence for gastrointestinal infection of SARS-CoV-2. *Gastroenterology*. March 2020. doi:10.1053/j.gastro.2020.02.055
3. Bai Y, Yao L, Wei T, et al. Presumed Asymptomatic Carrier Transmission of COVID-19. *JAMA*. February 2020. doi:10.1001/jama.2020.2565
4. van Doremalen N, Bushmaker T, Morris DH, et al. Aerosol and Surface Stability of SARS-CoV-2 as Compared with SARS-CoV-1. *New England Journal of Medicine*. 2020;0(0):null. doi:10.1056/NEJMc2004973
5. Li R, Pei S, Chen B, et al. Substantial undocumented infection facilitates the rapid dissemination of novel coronavirus (SARS-CoV2). *Science*. March 2020. doi:10.1126/science.abb3221
6. Wang D, Hu B, Hu C, et al. Clinical Characteristics of 138 Hospitalized Patients With 2019 Novel Coronavirus–Infected Pneumonia in Wuhan, China. *JAMA*. 2020;323:1061-1069.
7. Zhang Y, Zhang X, Liu L, Wang H, Zhao Q. Suggestions of Infection Prevention and Control in Digestive Endoscopy During Current 2019-nCoV Pneumonia Outbreak in Wuhan, Hubei Province, China. February 2020. <http://www.worldendo.org/wp-content/uploads/2020/02/Suggestions-of-Infection-Prevention-and-Control-in-Digestive-Endoscopy-During-Current-2019-nCoV-Pneumonia-Outbreak-in-Wuhan-Hubei-Province-China.pdf>. Accessed March 17, 2020.
8. Repici A, Maselli R, Colombo M, et al. Coronavirus (COVID-19) outbreak: what the department of endoscopy should know. *Gastrointestinal Endoscopy*. March 2020. doi:10.1016/j.gie.2020.03.019
9. Muscarella LF. Recommendations for the prevention of transmission of SARS during GI endoscopy. *Gastrointestinal Endoscopy*. 2004;60:792-795.
10. Pan L, Mu M, Ren HG, et al. Clinical characteristics of COVID-19 patients with digestive symptoms in Hubei, China: a descriptive, cross-sectional, multicenter study. :23.
11. *10 Things You Can Do to Manage COVID-19 at Home*. CDC; 2020. <https://www.youtube.com/watch?v=qPoptbtBjkg>.
12. Chapman S. Hot air? *BMJ*. 2001;323:1449.
13. Johnston ER, Habib-Bein N, Dueker JM, et al. Risk of bacterial exposure to the endoscopist's face during endoscopy. *Gastrointest Endosc*. 2019;89:818-824.
14. World Health Organization. *Rational Use of Personal Protective Equipment for Coronavirus Disease (COVID-19): Interim Guidance, 27 February 2020*. Geneva: World Health Organization; 2020. <https://extranet.who.int/iris/restricted/handle/10665/331215>.
15. Ong SWX, Tan YK, Chia PY, et al. Air, Surface Environmental, and Personal Protective Equipment Contamination by Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) From a Symptomatic Patient. *JAMA*. March 2020. doi:10.1001/jama.2020.3227
16. Knowlton SD, Boles CL, Perencevich EN, Diekema DJ, Nonnenmann MW, CDC Epicenters Program. Bioaerosol concentrations generated from toilet flushing in a hospital-based patient care setting. *Antimicrob Resist Infect Control*. 2018;7:16.
17. Adams JG, Walls RM. Supporting the Health Care Workforce During the COVID-19 Global Epidemic. *JAMA*. March 2020. doi:10.1001/jama.2020.3972
18. Longtin Y, Sax H, Allegranzi B, Schneider F, Pittet D. Hand Hygiene. *New England Journal of Medicine*. 2011;364:e24.
19. PPE | Appendix A | Isolation Precautions | Guidelines Library | Infection Control | CDC. <https://www.cdc.gov/infectioncontrol/guidelines/isolation/appendix/ppe.html>. Published March 26, 2019. Accessed March 17, 2020.
20. Ortega R, Bhadelia N, Obanor O, et al. Putting On and Removing Personal Protective

Equipment. *New England Journal of Medicine*. 2015;372(12):e16.  
doi:10.1056/NEJMvcm1412105

21. Implementation of Mitigation Strategies for Communities with Local COVID-19 Transmission. March 2020. <https://www.cdc.gov/coronavirus/2019-ncov/downloads/community-mitigation-strategy.pdf>. Accessed March 18, 2020.
22. Woelfel R, Corman VM, Guggemos W, et al. *Clinical Presentation and Virological Assessment of Hospitalized Cases of Coronavirus Disease 2019 in a Travel-Associated Transmission Cluster*. *Infectious Diseases (except HIV/AIDS)*; 2020. doi:10.1101/2020.03.05.20030502
23. van Doremalen N, Bushmaker T, Morris D, et al. *Aerosol and Surface Stability of HCoV-19 (SARS-CoV-2) Compared to SARS-CoV-1*. *Infectious Diseases (except HIV/AIDS)*; 2020. doi:10.1101/2020.03.09.20033217
24. CDC 2019-Novel Coronavirus (2019-nCoV) Real-Time RT-PCR Diagnostic Panel. Instructions for Use. March 2020. <https://www.fda.gov/media/134922/download>. Accessed March 18, 2020.
25. JOINT GI SOCIETY MESSAGE: COVID-19 Clinical Insights for Our Community of Gastroenterologists and Gastroenterology Care Providers. ASGE. <https://www.asge.org/home/joint-gi-society-message-covid-19>. Accessed March 18, 2020.



## Journal CME Conflict of Interest: Disclosure and Attestation

**Lead Author:** Roy M Soetikno

**Article:** Considerations in Performing Endoscopy During the COVID-19 Pandemic

**Date:** 03/17/2020

The purpose of this form is to identify all potential conflicts of interests that arise from financial relationships between any author for this article and any commercial or proprietary entity that produces healthcare-related products and/or services relevant to the content of the article. This includes any financial relationship within the last twelve months, as well as known financial relationships of authors' spouse or partner. **The lead author is responsible for submitting the disclosures of all listed authors, and must sign this form at the bottom. Additional forms may be submitted if the number of authors exceeds the space provided.**

**Lead Author: Roy M Soetikno MD MS**

**Email Address\*:** soetikno@earthlink.net

No financial relationships with a commercial entity producing health-care related products and/or services relevant to this article.

Company	Type of Relationship**	Content Area (if applicable)
Olympus	Consultant	
Fujifilm	Consultant	

**Author: Anthony YB Teoh MBChB**

**Email Address\*:** anthonyteoh@surgery.cuhk.edu.hk

No financial relationships with a commercial entity producing health-care related products and/or services relevant to this article.

Company	Type of Relationship**	Content Area (if applicable)
Boston Scientific	Consultant	
Cook	Consultant	
Taewoong and Microtech	Consultant	

**Author: Ravishankar Asokkumar MBBS**

**Email Address\*:**  
ravishankar.asokkumar@singhealth.com.sg

No financial relationships with a commercial entity producing health-care related products and/or services relevant to this article.

Company	Type of Relationship**	Content Area (if applicable)

**Author: Patricia Cabral-Prodigalidad MD**

**Email Address\*:** scprodigalidad@yahoo.com

No financial relationships with a commercial entity producing health-care related products and/or services relevant to this article.

Company	Type of Relationship**	Content Area (if applicable)

\* We will use email addresses only for questions related to this article 1

\*\* **Type of relationship may include:** full-time or part-time employee, independent contractor, consultant, research or other grant recipient, paid speaker or teacher, membership on advisory committee or review panels, ownership interest (product royalty/licensing fees, owning stocks, shares, etc.), relationship of a spouse or partner, or any other financial relationship.





**Author: Tonya Kaltenbach MD MS**

**Email Address\*: endoresection@me.com**

No financial relationships with a commercial entity producing health-care related products and/or services relevant to this article.

Company	Type of Relationship**	Content Area (if applicable)
Olympus	Consultant	
Aries Pharmaceuticals	Consultant	
Medtronic	Consultant	

**Author: James YW Lau MD**

**Email Address\*: laujyw@surgery.cuhk.edu.hk**

No financial relationships with a commercial entity producing health-care related products and/or services relevant to this article.

Company	Type of Relationship**	Content Area (if applicable)

**Author: Amandeep Shergill MD MS**

**Email Address\*: amandeep.shergill@ucsf.edu**

No financial relationships with a commercial entity producing health-care related products and/or services relevant to this article.

Company	Type of Relationship**	Content Area (if applicable)
Pentax	Research gift recipient	

**Author:**

**Email Address\*:**

No financial relationships with a commercial entity producing health-care related products and/or services relevant to this article.

Company	Type of Relationship**	Content Area (if applicable)

**As corresponding author of this article**, I attest that I have received disclosure information from all participating authors as listed above and acknowledge that I am responsible for verifying the accuracy of and reporting completely the information provided to me. Financial relationships relevant to this article can be researched at <https://www.cms.gov/openpayments/>. I understand that typing my name below serves as an electronic signature for the purposes of this form.

***Roy M Soetikno***

**Type Name (Electronic Signature)**

\* We will use email addresses only for questions related to this article 2

\*\* **Type of relationship may include:** full-time or part-time employee, independent contractor, consultant, research or other grant recipient, paid speaker or teacher, membership on advisory committee or review panels, ownership interest (product royalty/licensing fees, owning stocks, shares, etc.), relationship of a spouse or partner, or any other financial relationship.

